

February 1, 2008

# PHASE SEPARATION AND THREE-SITE HOPPING IN THE 2-DIMENSIONAL $t$ - $J$ MODEL

Elisa Ercolessi<sup>a)</sup>, Pierbiagio Pieri<sup>a,b)</sup>, Marco Roncaglia<sup>a,b)</sup>

<sup>a)</sup>*Dipartimento di Fisica, Università di Bologna and INFN  
Via Irnerio 46, I-40126, Bologna, Italy.*  
<sup>b)</sup>*INFN, Sezione di Bologna, Italy.*

## Abstract

We study the  $t$ - $J$  model with the inclusion of the so called three-site term which comes out from the  $t/U \rightarrow 0$  expansion of the Hubbard model. We find that this singlet pair hopping term has no qualitative effect on the structure of the pure mean field phase diagram for nonmagnetic states. In accordance with experimental data on high- $T_c$  materials and some numerical studies, we also find wide regions of phase coexistence whenever the coupling  $J$  is greater than a critical value  $J_c$ . We show that  $J_c$  varies linearly with the temperature  $T$ , going to zero at  $T = 0$ .

PACS: 71.27, 74.20.

In a recent paper [1] we have presented a new finite temperature phase diagram for the nonmagnetic phases (dimer and flux) in the 2-dimensional  $t$ - $J$  model below half-filling, which shows the existence of broad areas of phase coexistence (dimer-flux or flux-uniform), in accordance with experimental and numerical data on the possibility of separation into hole-rich and hole-poor regions [2, 3, 4].

It is well known that one can obtain the  $t$ - $J$  model in the strong coupling regime ( $U \gg t$ ) of the one-band Hubbard model:

$$H = -t \sum_{\langle ij \rangle} \sum_{\sigma} c_{i\sigma}^{\dagger} c_{j\sigma} + U \sum_i n_{i\uparrow} n_{i\downarrow} \quad (1)$$

below half-filling. In (1) Latin indices label sites, the  $c_{i\sigma}$ 's are annihilation operators for a fermion on site  $i$  and spin  $\sigma = \uparrow, \downarrow$ ,  $n_{i\sigma} = c_{i\sigma}^{\dagger} c_{i\sigma}$  and  $\langle ij \rangle$  denotes a sum over ordered nearest neighbor (n.n.) pairs.

The strong coupling limit of (1) can be studied either by direct perturbation methods [5] or by canonical transformations methods [6] (see also [7, 8] for a detailed review). To lowest order in  $t/U$ , the Hamiltonian that describes the Hubbard model on the subspace with no double occupancies ( $n_i \leq 1$ ) is given by:

$$\widehat{H} = T_h + H^{(1)} + H^{(2)} \quad (2)$$

where

$$T_h = - \sum_{\langle ij \rangle \sigma} t_{ij} (1 - n_{i\bar{\sigma}}) c_{i\sigma}^{\dagger} c_{j\sigma} (1 - n_{j\bar{\sigma}}) ; \bar{\sigma} = -\sigma \quad (3)$$

$$H^{(1)} = \sum_{\langle ij \rangle} J_{ij} \left\{ \vec{S}_i \cdot \vec{S}_j - \frac{1}{4} n_i n_j \right\} , \quad J_{ij} = 4 \frac{|t_{ij}|^2}{U} , \quad (4)$$

$$H^{(2)} = -\frac{1}{U} \sum_{\langle ijl \rangle} \sum_{\sigma\tau} t_{ij} t_{jl} (1 - n_{i\bar{\sigma}}) c_{i\sigma}^{\dagger} c_{j\sigma} n_{j\bar{\sigma}} n_{j\bar{\tau}} c_{j\tau}^{\dagger} c_{l\tau} (1 - n_{l\bar{\tau}}) \quad (5)$$

and the symbol  $\langle ij \rangle$  denotes a summation over unordered pairs of n.n. and  $\langle ijl \rangle$  denotes a summation where  $i \neq l$ , and both  $i$  and  $l$  are n.n. to  $j$ . The spin operators are defined as

$$\vec{S}_i = \frac{1}{2} \sum_{\alpha\beta} c_{i\alpha}^{\dagger} \vec{\sigma}_{\alpha\beta} c_{i\beta} , \quad (6)$$

with  $\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$  being Pauli matrices.

The effective Hamiltonian contains a direct hopping term  $T_h$  that gives nonvanishing contribution only if the electron jumps from site  $i$  to an empty n.n. site  $j$ . So we expect the effective hopping coefficient  $t$  to be reduced by a factor of order  $\delta$ ,  $\delta$  being the doping fraction [9]. On the contrary  $H^{(1)}$ , which gives rise to a spin dynamics described by the standard AFM Heisenberg model, has not to be renormalized, since  $\vec{S}_i \cdot \vec{S}_j$  is nonzero only if both the sites  $i$  and  $j$  are singly occupied.  $H^{(2)}$  is proportional to  $|t|^2$  and describes a double hopping process. Thus, one would be tempted to believe that it has to be

renormalized by a factor of  $\delta^2$ . However, it is not difficult to see that the three-site hopping term  $H^{(2)}$  gives nonzero contribution only for the situation in which the final site  $l$  is empty and the intermediate site  $j$  is singly occupied. Then this term should be renormalized by a factor of  $\delta(1 - \delta)$ . Similar conclusions can be reached within different approaches [10, 11, 12].

For low doping the three-site term is smaller by a factor  $\delta$  as compared with the AFM term and by a factor  $|t|/U$  as compared with the direct hopping term. That is why it is usually ignored and one is led to the  $t$ - $J$  model [13]. However, the Hubbard Hamiltonian is often studied in a range of the parameter  $t/U$  not so small, especially in applications to high  $T_c$  materials for which one assumes  $t/U \approx 0.1$ . In this regime the role of the three-site term cannot be neglected a priori.

Numerical works have recently shown that the three-site term is essential to avoid discrepancies between the one-particle spectra of the Hubbard and  $t$ - $J$  model in the intermediate  $U$  regime [14]. According to [14], only after the inclusion of the three-site term, the  $t$ - $J$  model can correctly describe the photoemission experiments on cuprate oxides.

The three-site term describes the hopping of a singlet pair, which can be either rigid or with a flip of the spins. Both these processes preserve the local singlet bond and may be relevant for superconducting correlations, favoring superconducting states at the expense of an antiferromagnetic phase [10].

Therefore, according to the discussion above, we approximate the effect of the Gutzwiller projector by renormalizing each term of the Hamiltonian (2) with suitable factors depending on  $\delta$ . Our new starting point is hence the Hamiltonian:

$$\mathcal{H} = t\delta \sum_{(ij)} \sum_{\alpha} c_{i\alpha}^{\dagger} c_{j\alpha} - \frac{J}{4} \sum_{ijkl} \sum_{\alpha\beta} c_{i\alpha}^{\dagger} c_{j\alpha} \mathcal{A}_{ji,kl} c_{l\beta}^{\dagger} c_{k\beta} \quad (7)$$

where we have introduced the four fermion matrix

$$\mathcal{A}_{ji,kl} = \begin{cases} 1 & \text{if } i \text{ n.n. } j \text{ and } i = k, j = l \\ \delta(1 - \delta) & \text{if } i \text{ n.n. } j, k \text{ n.n. } l, j = l \text{ and } i \neq k \\ 0 & \text{otherwise} \end{cases} \quad (8)$$

In [1], we examined the so-called nonmagnetic phases for the  $t$ - $J$  model, first proposed by Affleck and Marston [15], in the mean field approximation and we have presented a finite temperature phase diagram ( $T$  versus the doping  $\delta$ ). In this paper we shall study how the phase diagram changes if we include the three-site hopping term.

To study the Hamiltonian (7) we can parallel the discussion in [1], where we have chosen to represent the partition function in the grand canonical ensemble by means of a functional integral over Grassmann fields and decoupled the four-fermion interaction by means of a Hubbard-Stratonovich transformations over the auxiliary fields  $\mathcal{U}_{ij}(\tau) = \mathcal{U}_{ji}^{\dagger}(\tau)$ . In the static approximation  $\mathcal{U}_{ij}(\tau) \equiv \mathcal{U}_{ij}$ , the partition function reduces to:

$$\mathcal{Z} = \int [\mathcal{D}\mathcal{U}_{ij}^* \mathcal{D}\mathcal{U}_{ij}] \exp \left\{ -\frac{\beta}{J} \sum_{ijkl} \mathcal{U}_{ij}^* \mathcal{A}_{ji,kl} \mathcal{U}_{kl} \right\} \exp \{ -\mathcal{S}_{eff} \} \quad (9)$$

$$\mathcal{S}_{eff} \equiv -2 \text{Tr} \left\{ \log \left[ -\mathbb{G}^{-1} + \beta \mathbb{U} \right] \right\}, \quad (10)$$

where

$$[\mathbb{G}^{-1}]_{nn'}^{ij} = (i\omega_n + \mu\beta) \delta_{ij} \delta_{nn'} \quad (11)$$

$$[\mathbb{U}]_{nn'}^{ij} = \begin{cases} [t\delta + \sum_{pq} \mathcal{U}_{pq} \mathcal{A}_{pq,ji}] \delta_{nn'} & \text{if } i \text{ n.n. } j \\ 0 & \text{otherwise} \end{cases}, \quad (12)$$

$\mu$  being the chemical potential.

We consider, as in [15], a pattern of link variables  $\mathcal{U}_{ij}$  that admits a symmetry for translations along the diagonal so that the matrix  $\mathbb{U}$  depends only on the four parameters  $\mathcal{U}_j$  ( $j = 1, \dots, 4$ ) shown in Fig. 1 of [1]. As for the pure  $t$ - $J$  model, the matrix  $\mathbb{U}$  can be easily diagonalized in momentum space: if we include the three-site term it has eigenvalues

$$\begin{aligned} E_k &= \pm |\lambda_k| \\ \lambda_k &\equiv \chi_1 e^{ik_x a} + \chi_2^* e^{-ik_y a} + \chi_3 e^{-ik_x a} + \chi_4^* e^{ik_y a} \end{aligned} \quad (13)$$

with the definitions

$$\chi_1 \equiv t\delta + \mathcal{U}_1 + \delta(1 - \delta)(\mathcal{U}_2^* + \mathcal{U}_3 + \mathcal{U}_4^*) \quad (14)$$

$$\chi_2 \equiv t\delta + \mathcal{U}_2 + \delta(1 - \delta)(\mathcal{U}_1^* + \mathcal{U}_3^* + \mathcal{U}_4) \quad (15)$$

$$\chi_3 \equiv t\delta + \mathcal{U}_3 + \delta(1 - \delta)(\mathcal{U}_1 + \mathcal{U}_2^* + \mathcal{U}_4^*) \quad (16)$$

$$\chi_4 \equiv t\delta + \mathcal{U}_4 + \delta(1 - \delta)(\mathcal{U}_1^* + \mathcal{U}_2 + \mathcal{U}_3^*) \quad (17)$$

We notice that both the expression for the partition function (9) and for the eigenvalues of  $\mathbb{U}$  (13) are formally identical to the one for the pure  $t$ - $J$  model we gave in [1], the only differences being in the definition of the matrix  $\mathcal{A}$  and of the parameters  $\chi_j$ 's that appear in  $\lambda_k$ . Thus, to study nonmagnetic phases in the mean field approximation, we can borrow from the discussion of [1].

We first discuss in detail the numerical solutions to the self-consistency equations for the case  $J/t = 1$ . For high temperature and/or high doping the free energy is minimized by the uniform phase, characterized by  $\mathcal{U}_i \equiv \mathcal{U} \in \mathbb{R}^+$ ,  $i = 1, \dots, 4$ . For temperature lower than  $T = J/4$  we find two phases: dimer and flux. The former corresponds to two solutions of the form  $\mathcal{U}_1 \neq \mathcal{U}_2 = \mathcal{U}_3 = \mathcal{U}_4$ ,  $\mathcal{U}_j \in \mathbb{C}$  with  $|\mathcal{U}_1| \gg |\mathcal{U}_2|$  and is stable in the lower range of dopings. In the flux phase the link variables  $\mathcal{U}_i$  are all equal to a complex value  $\mathcal{U}e^{i\phi}$  with  $\phi \neq 0$ . It is the lowest energy state in a range of doping that roughly




Figure 1:  
Mean field phase diagram for the  $t$ - $J$  model (open circles)  
and with the inclusion of the three-site term (solid triangles).

corresponds to that of high- $T_c$  superconductivity. The pure mean field diagram is shown in Fig. 1 (solid triangles).

Since the hole mobility is enhanced by the addition of the three-site hopping term, we expect disordered phases be favored with respect to the ordered ones. Moreover, the three-site term being proportional to  $\delta(1 - \delta)$ , this effect should increase with doping in the region we considered. This is in fact what we find in our mean field calculation. Fig. 1 shows the shifts of the phase boundaries for the usual  $t$ - $J$  model (open circles) after the inclusion of the three-site term (solid triangles). The region of stability of the dimer phase shrinks in favor of flux phase, while the uniform configuration extends to considerably lower dopings at the expense of the flux one.

As we have observed also in [1], for  $T = 0$  our mean field calculations are consistent with the ones carried out within a slave boson formalism in [16, 17]. In particular, we agree with [17] in finding a first order phase transition between the dimer and the flux phases. Contrary to [17], we find that the flux-uniform phase transition is of the second order for any temperature. This can be seen from either Fig. 3 or Fig. 5 of [1]. In Fig. 3 of [1] it is possible to observe that, increasing  $\delta$ , the angle  $\phi$ , which distinguishes the flux phase from the uniform, decreases continuously from  $\pi/4$  to 0. In Fig. 5 of [1], it is easy to see that, at the transition point, there is no jump in the chemical potential, but

only a cusp. At this regard, we can also observe that the slope of these curves becomes steeper as the temperature decreases. For  $T = 0$  this behaviour gives rise to numerical problems and probably this is the reason why in [17] the authors found a jump in the chemical potential and hence concluded that the flux-uniform boundary line describes a first order phase transition.

The diagram of Fig. 1 is constructed without taking into account the possibility of coexistence of different phases. Indeed we find two ranges, belonging respectively to the dimer and the flux phase, where  $\frac{\partial \mu}{\partial \delta} > 0$  [1, 18] which clearly indicate an instability towards phase separation. By means of the Maxwell construction on the chemical potential [4], we obtain two different regions corresponding to dimer-flux and flux-uniform phase coexistence. In the former, the hole-poor (hole-rich) part is in the dimer (flux) phase while in the latter the hole-poor (hole-rich) part corresponds to the flux (uniform) phase. The final phase diagram is shown in Fig. 2.




Figure 2:

The phase diagram after taking into account phase separation.

Phase separation for nonmagnetic phases is also discussed in [17]. In the range of dopings relevant to high- $T_c$  superconductivity, the authors of [17] found a wide region of phase separation between a half-filling dimer phase and a hole-rich *uniform* phase, while we find a smaller region of phase separation between a half-filling dimer phase and a *flux* phase corresponding to  $\delta \approx 0.09$ . This discrepancy is due to a different behaviour of the chemical potential as a function of  $\delta$ . Indeed, within the large- $N$  slave-boson formalism the

electronic chemical potential gets shifted by the mean value of the Lagrange multiplier that enforces the constraint, while in the approach we adopted such a term is absent. Phase separation is a relevant issue because there is a considerable experimental evidence that many high- $T_c$  superconductors have a regime in which phase separation between an AFM hole-poor region and a superconducting hole-rich region occurs [19]. Also, according to [20] an instability toward phase separation might explain the unusual properties of the normal state and superconductivity in copper oxides. This is why the above mentioned discrepancies between different approaches deserve a deeper investigation, which we plan to describe in a future work.




Figure 3:  
The region of antiferromagnetic couplings where phase separation occurs.  
The dotted line refers to a calculation without the three-site term.

We want finally to give our contribution to the open question about the possibility that phase separation disappears once  $J/t$  is lowered below a critical value  $J_c/t$ . The determination of this quantity is important in order to understand if the  $t$ - $J$  model can exhibit phase separation in the regime of high- $T_c$  superconductors where  $0.3 < J/t < 0.5$ . Theoretical studies, for  $T = 0$ , based on different techniques, lead to contradictory results. Using a high temperature expansion the authors of [18] find the model phase separates only for  $J/t > 1$ , i.e. outside the region interesting for superconductivity. Similar results have been obtained in some numerical works [21]. On the contrary other authors find that phase separation occurs for all values of  $J/t$ . In particular this has been done by means of quantum Monte Carlo calculations [22], a combination of numerical

and analytical results [2, 3] and the above mentioned slave-boson approach [4, 17]. We calculated numerically the critical value  $J_c/t$  for several temperatures ( $T = 0$  included) by observing the disappearance of the instability regions where  $\frac{\partial \mu}{\partial \phi} > 0$ .

The results, plotted in Fig. 3, show that  $J_c/t$  varies linearly with the temperature and vanishes for  $T = 0$ , both for the standard  $t$ - $J$  model (dotted line) and with the inclusion of the three-site term (solid line). Thus, at zero temperature we find phase separation for all values of the interaction strength, in accordance with [3, 17, 22]. In addition, we can also conclude that at finite temperature there is always a nonzero critical value  $J_c/t$  below which phase separation disappears. In particular there is no phase separation in the region interesting for superconductivity for temperatures higher than  $T \approx 0.07t$ .

Lastly, we observe that in one dimension some exact diagonalization and Monte Carlo results [23] suggest that the inclusion of the three-site term suppresses phase separation, which is instead present when such a term is not added [24, 25]. This seems not to be the case in two dimensions, at least at mean field level. Indeed, our calculations show that phase separation is still present and the effect of the three-site term is not so dramatic.

### Acknowledgments

We are indebted to G. Morandi for many valuable suggestions. We are also grateful to D. Baeriswyl, F. Napoli and G. C. Strinati for helpful discussions. Finally, we would like to thank the referee for suggesting us to look for the occurrence of phase separation also at small values of  $J/t$ .



## References

- [1] E. Ercolessi, P. Pieri and M. Roncaglia, Phys. Lett. **A225** (1997) 331-340
- [2] E. Dagotto and J. Riera, Phys. Rev. Lett. **70** (1993) 682;  
A. Angelucci and S. Sorella, Phys. Rev. **B47** (1993) 8858;  
M. Marder, N. Papanicolaou and G.C. Psaltakis, Phys. Rev. **B41** (1990) 6920;  
A. Auerbach, B. E. Larson, Phys. Rev. B **43** (1991) 7800;
- [3] V.J. Emery, S.A. Kivelson and H.Q. Lin, Phys. Rev. Lett. **64** (1990) 475;  
S.A. Kivelson, V.J. Emery and H.Q. Lin, Phys. Rev. **B42** (1990) 6523
- [4] M. Grilli, R. Raimondi, C. Castellani, C. Di Castro, G. Kotliar, Phys. Rev. Lett. **67** (1991) 259
- [5] V.J. Emery, Phys. Rev. **B14** (1976) 2989;
- [6] M. Aquarone, “Hubbard Correlations in a Single Band”, in *Physics of Metals*, eds. E.S. Giuliano and C. Rizzuto, (World Scientific, 1988) and references therein;  
K.A. Chao, J. Spalek and A.M. Oles, J. Phys. **C10** (1977) L271;  
K.A. Chao, J. Spalek and A.M. Oles, Phys. Rev. **B18** (1978) 3453;  
C. Gros, R. Joynt and T.M. Rice, Phys. Rev. **B36** (1987) 381;  
H. Primas, Helv. Phys. Acta **34** (1961) 331;  
H. Primas, Rev. Mod. Phys. **35** (1963) 710
- [7] A.P. Balachandran, E. Ercolessi, G. Morandi and A.M. Srivastava, *Hubbard Model and Anyon Superconductivity*, World Scientific (1990)
- [8] E. Galleani d’Agliano, G. Morandi and F. Napoli, in *High Temperature Superconductivity*, M. Acquarone Ed., World Scientific (1996)
- [9] G. Baskaran, Z. Zou and P.W. Anderson, Solid State Comm. **63** (1987) 973
- [10] J. Spalek, Phys. Rev. **B37** (1988) 533
- [11] G. Kotliar and J. Liu, Phys. Rev. **B38** (1988) 5142
- [12] M. Grilli and G. Kotliar, Phys. Rev. Lett. **64** (1990) 1170
- [13] J.E. Hirsch, Phys. Rev. Lett. **54** (1985) 1317;  
C. Gros, R. Joynt and T.M. Rice, Phys. Rev. **B36** (1987) 8190
- [14] H. Eskes and R. Eder, Phys. Rev. **B54** (1996) R14226
- [15] J.B. Marston and I. Affleck, Phys. Rev. **B39** (1989) 11538 ;  
I. Affleck and J.B. Marston, Phys. Rev. **B37** (1988) 3774

- [16] M.U. Ubbens and P.A. Lee, Phys. Rev. **B46** (1992) 8434;  
X.G. Wen and P.A. Lee, Phys. Rev. Lett. **76** (1996) 503;  
D.N. Sheng, Z.B. Su and L. Yu, Phys. Rev. **B42** (1990) 8732
- [17] M. Grilli, C. Castellani and G. Kotliar, Phys. Rev. **B45** (1992) 10805
- [18] W. O. Putikka, M. U. Luchini, T. M. Rice, Phys. Rev. Lett. **68** (1992) 538
- [19] M.F. Hundley, J.D. Thompson, S.W. Cheong, Z. Fisk and J.E. Schirber, Phys. Rev. **B41** (1990) 4062;  
J.H. Cho, F.C. Chou and D.C. Johnston, Phys. Rev. Lett. **70** (1993) 222;  
J.M. Tranquada, B.J. Sternlieb, D.J. Axe, Y. Nakamura and S. Uchida, Nature **375** (1995) 561
- [20] A. Perali, C. Castellani, C. Di Castro and M. Grilli, Phys. Rev. **B54** (1996) 16216
- [21] P. Prelovsek and X. Zotos, Phys. Rev. **B47** (1993) 5984;  
D. Poilblanc, Phys. Rev. **B52** (1995) 9201
- [22] C. S. Hellberg and E. Manousakis, cond-mat/9611195
- [23] B. Ammon, M. Troyer and H. Tsunetsugu, Phys. Rev. **B52** (1995) 629
- [24] Masao Ogata, M. U. Luchini, S. Sorella and F. F. Assaad, Phys. Rev. Lett. **66** (1991) 2388
- [25] F. F. Assaad and D. Würtz, Phys. Rev. **B44** (1991) 2681